arXiv:1511.07318v1 [stat.ML] 20 Nov 2015

Bayesian SPLDA

Jesús Villalba

Communications Technology Group (GTC),
Aragon Institute for Engineering Research (I3A),
University of Zaragoza, Spain
villalba@unizar.es

May 15, 2012

1 Introduction

In this document we are going to derive the equations needed to implement a Variational Bayes estimation of the parameters of the SPLDA model [1]. This can be used to adapt the SPLDA from one database to another with few development data or to implement the fully Bayesian recipe [2]. Our approach is similar to Bishop's VB PPCA in [3].

2 The Model

2.1 SPLDA

SPLDA is a linear generative model represented in Figure 1.



Figure 1: BN for Bayesian SPLDA model.

An i-vector ϕ of speaker i can be written as:

$$\phi_{ij} = \mu + \mathbf{V}\mathbf{y}_i + \epsilon_{ij} \tag{1}$$

where μ is a speaker independent mean, $\mathbf{V}$ is the eigenvoices matrix, $\mathbf{y}_i$ is the speaker factor vector, and ϵ is a channel offset.

We assume the following priors for the variables:

$$\mathbf{y}_i \sim \mathcal{N}(\mathbf{y}_i|\mathbf{0}, \mathbf{I}) \tag{2}$$

$$\epsilon_{ij} \sim \mathcal{N}\left(\epsilon_{ij}|\mathbf{0}, \mathbf{W}^{-1}\right) \tag{3}$$

where $\mathcal{N}$ denotes a Gaussian distribution; $\mathbf{W}$ is the within class precision matrix.

2.2 Notation

We are going to introduce some notation:

- Let $\mathbf{\Phi}_\mathrm{d}$ be the development i-vectors dataset.
- Let $\mathbf{\Phi}_\mathrm{t} = \{l, r\}$ be the test i-vectors.
- Let $\mathbf{\Phi}$ be any of the previous datasets.
- Let θ_d be the labelling of the development dataset. It partitions the N_d i-vectors into M_d speakers.
- Let θ_t be the labelling of the test set, so that $\theta_\mathrm{t} \in \{\mathcal{T}, \mathcal{N}\}$, where $\mathcal{T}$ is the hypothesis that l and r belong to the same speaker and $\mathcal{N}$ is the hypothesis that they belong to different speakers.
- Let θ be any of the previous labellings.
- Let $\mathbf{\Phi}_i$ be the i-vectors belonging to the speaker i.
- Let ϕ_{ij} the i-vector j of speaker i.
- Let $\mathbf{Y}_\mathrm{d}$ be the speaker identity variables of the development set. We will have as many identity variables as speakers.
- Let $\mathbf{Y}_\mathrm{t}$ be the speaker identity variables of the test set.
- Let $\mathbf{Y}$ be any of the previous speaker identity variables sets.
- Let d be the i-vector dimension.
- Let n_y be the speaker factor dimension.
- Let $\mathcal{M} = (\mu, \mathbf{V}, \mathbf{W})$ be the set of all the parameters.

3 Sufficient statistics

We define the sufficient statistics for speaker i. The zero-order statistic is the number of observations of speaker i N_i. The first-order and second-order statistics are

$$\mathbf{F}_i = \sum_{j=1}^{N_i} \phi_{ij} \tag{4}$$

$$\mathbf{S}_i = \sum_{j=1}^{N_i} \phi_{ij}\phi_{ij}^T \tag{5}$$

We define the centered statistics as

$$\overline{\mathbf{F}}_i = \mathbf{F}_i - N_i\mu \tag{6}$$

$$\overline{\mathbf{S}}_i = \sum_{j=1}^{N_i} (\phi_{ij} - \mu)(\phi_{ij} - \mu)^T = \mathbf{S}_i - \mu\mathbf{F}_i^T - \mathbf{F}_i\mu^T + N_i\mu\mu^T \tag{7}$$

We define the global statistics

$$N = \sum_{i=1}^{M} N_i \tag{8}$$

$$\mathbf{F} = \sum_{i=1}^{M} \mathbf{F}_i \tag{9}$$

$$\overline{\mathbf{F}} = \sum_{i=1}^{M} \overline{\mathbf{F}}_i \tag{10}$$

$$\mathbf{S} = \sum_{i=1}^{M} \mathbf{S}_i \tag{11}$$

$$\overline{\mathbf{S}} = \sum_{i=1}^{M} \overline{\mathbf{S}}_i \tag{12}$$

4 Data conditional likelihood

The likelihood of the data given the hidden variables for speaker i is

$$\ln P\left(\mathbf{\Phi}_i|\mathbf{y}_i, \mu, \mathbf{V}, \mathbf{W}\right) = \sum_{j=1}^{N_i} \ln \mathcal{N}\left(\phi_{ij}|\mu + \mathbf{V}\mathbf{y}_i, \mathbf{W}^{-1}\right) \tag{13}$$

$$= \frac{N_i}{2} \ln \left|\frac{\mathbf{W}}{2\pi}\right| - \frac{1}{2} \sum_{j=1}^{N_i} (\phi_{ij} - \mu - \mathbf{V}\mathbf{y}_i)^T \mathbf{W} (\phi_{ij} - \mu - \mathbf{V}\mathbf{y}_i) \tag{14}$$

$$= \frac{N_i}{2} \ln \left|\frac{\mathbf{W}}{2\pi}\right| - \frac{1}{2} \mathrm{tr}\left(\mathbf{W}\overline{\mathbf{S}}_i\right) + \mathbf{y}_i^T \mathbf{V}^T \mathbf{W} \overline{\mathbf{F}}_i - \frac{N_i}{2} \mathbf{y}_i^T \mathbf{V}^T \mathbf{W} \mathbf{V} \mathbf{y}_i \tag{15}$$

We can write this likelihood in another form:

$$\ln P\left(\mathbf{\Phi}_i|\mathbf{y}_i, \mu, \mathbf{V}, \mathbf{W}\right) = \frac{N_i}{2} \ln \left|\frac{\mathbf{W}}{2\pi}\right| - \frac{1}{2} \mathrm{tr}\left(\mathbf{W}\left(\mathbf{S}_i - 2\mathbf{F}_i \mu^T + N_i \mu \mu^T\right.\right. \tag{16}$$

$$\left.\left. -2\left(\mathbf{F}_i - N_i \mu\right) \mathbf{y}_i^T \mathbf{V}^T + N_i \mathbf{V} \mathbf{y}_i \mathbf{y}_i^T \mathbf{V}^T\right)\right) \tag{17}$$

We can write this likelihood in another form if we define:

$$\tilde{\mathbf{y}}_i = \begin{bmatrix} \mathbf{y}_i \\ 1 \end{bmatrix}, \quad \tilde{\mathbf{V}} = \begin{bmatrix} \mathbf{V} & \mu \end{bmatrix} \tag{18}$$

Then

$$\ln P\left(\mathbf{\Phi}_i|\mathbf{y}_i, \mu, \mathbf{V}, \mathbf{W}\right) = \sum_{j=1}^{N_i} \ln \mathcal{N}\left(\phi_{ij}|\tilde{\mathbf{V}}\tilde{\mathbf{y}}_i, \mathbf{W}^{-1}\right) \tag{19}$$

$$= \frac{N_i}{2} \ln \left|\frac{\mathbf{W}}{2\pi}\right| - \frac{1}{2} \sum_{j=1}^{N_i} (\phi_{ij} - \tilde{\mathbf{V}}\tilde{\mathbf{y}}_i)^T \mathbf{W} (\phi_{ij} - \tilde{\mathbf{V}}\tilde{\mathbf{y}}_i) \tag{20}$$

$$= \frac{N_i}{2} \ln \left|\frac{\mathbf{W}}{2\pi}\right| - \frac{1}{2} \mathrm{tr}\left(\mathbf{W}\mathbf{S}_i\right) + \tilde{\mathbf{y}}_i^T \tilde{\mathbf{V}}^T \mathbf{W} \mathbf{F}_i - \frac{N_i}{2} \tilde{\mathbf{y}}_i^T \tilde{\mathbf{V}}^T \mathbf{W} \tilde{\mathbf{V}} \tilde{\mathbf{y}}_i \tag{21}$$

$$= \frac{N_i}{2} \ln \left|\frac{\mathbf{W}}{2\pi}\right| - \frac{1}{2} \mathrm{tr}\left(\mathbf{W}\left(\mathbf{S}_i - 2\mathbf{F}_i \tilde{\mathbf{y}}_i^T \tilde{\mathbf{V}}^T + N_i \tilde{\mathbf{V}} \tilde{\mathbf{y}}_i \tilde{\mathbf{y}}_i^T \tilde{\mathbf{V}}^T\right)\right) \tag{22}$$

5 Variational inference with Gaussian-Gamma priors for V, Gaussian for μ and Wishart for W (informative and non-informative)

5.1 Model priors

We introduce a *hierarchical* prior $P(\mathbf{V}|\alpha)$ over the matrix $\mathbf{V}$ governed by a n_y dimensional vector of hyperparameters where n_y is the dimension of the factors. Each hyperparameter controls one of the columns of the matrix $\mathbf{V}$ through a conditional Gaussian distribution of the form:

$$P(\mathbf{V}|\alpha) = \prod_{q=1}^{n_y} \left(\frac{\alpha_q}{2\pi}\right)^{d/2} \exp\left(-\frac{1}{2}\alpha_q \mathbf{v}_q^T \mathbf{v}_q\right) \tag{23}$$

where $\mathbf{v}_q$ are the columns of $\mathbf{V}$. Each α_q controls the inverse variance of the corresponding $\mathbf{v}_q$. If a particular α_q has a posterior distribution concentrated at large values, the corresponding $\mathbf{v}_q$ will tend to be small, and that direction of the latent space will be effectively 'switched off'.

We define a prior for α:

$$P(\alpha) = \prod_{q=1}^{n_y} \mathcal{G}(\alpha_q|a_\alpha, b_\alpha) \tag{24}$$

where $\mathcal{G}$ denotes the Gamma distribution. Bishop defines broad priors setting $a = b = 10^{-3}$.

We place a Gaussian prior for the mean μ:

$$P(\mu) = \mathcal{N}\left(\mu|\mu_0, \mathrm{diag}(\beta)^{-1}\right) \tag{25}$$

We will consider the case where each dimension has different precision and the case with isotropic precision ($\mathrm{diag}(\beta) = \beta\mathbf{I}$).

Finally, we put a Wishart prior on $\mathbf{W}$,

$$P(\mathbf{W}) = \mathcal{W}(\mathbf{W}|\mathbf{\Psi}_0, \nu_0) \tag{26}$$

We can make the Wishart prior non-informative like in [4].

$$P(\mathbf{W}) = \lim_{k\to 0} \mathcal{W}(\mathbf{W}|\mathbf{W}_0/k, k) \tag{27}$$

$$= \alpha\,|\mathbf{W}|^{-(d+1)/2} \tag{28}$$

5.2 Variational distributions

We write the joint distribution of the observed and latent variables:

$$P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha|\mu_0, \beta, a_\alpha, b_\alpha) = P(\mathbf{\Phi}|\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W})\, P(\mathbf{Y})\, P(\mathbf{V}|\alpha)\, P(\alpha|a, b)\, P(\mu|\mu_0, \beta)\, P(\mathbf{W}) \tag{29}$$

Following, the conditioning on $(\mu_0, \beta, a_\alpha, b_\alpha)$ will be dropped for convenience.

Now, we consider the partition of the posterior:

$$P(\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha|\mathbf{\Phi}) \approx q(\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha) = q(\mathbf{Y}) \prod_{r=1}^{d} q(\tilde{\mathbf{v}}_r')\, q(\mathbf{W})\, q(\alpha) \tag{30}$$

where $\tilde{\mathbf{v}}_r'$ is a column vector containing the r^{th} row of $\tilde{\mathbf{V}}$. If $\mathbf{W}$ were a diagonal matrix the factorization $\prod_{r=1}^{d} q(\tilde{\mathbf{v}}_r')$ is not necessary because it arises naturally when solving the posterior. However, for full covariance $\mathbf{W}$, the posterior of $\mathrm{vec}(\tilde{\mathbf{V}})$ is a Gaussian with a huge full covariance matrix. Therefore, we force the factorization to made the problem tractable.

The optimum for $q^*(\mathbf{Y})$:

$$\ln q^*(\mathbf{Y}) = \mathrm{E}_{\mu,\mathbf{V},\mathbf{W},\alpha}\left[\ln P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha)\right] + \mathrm{const} \tag{31}$$

$$= \mathrm{E}_{\mu,\mathbf{V},\mathbf{W}}\left[\ln P(\mathbf{\Phi}|\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W})\right] + \ln P(\mathbf{Y}) + \mathrm{const} \tag{32}$$

$$= \sum_{i=1}^{M} \mathbf{y}_i^T \mathrm{E}\left[\mathbf{V}^T\mathbf{W}(\mathbf{F}_i - N_i\mu)\right] - \frac{1}{2}\mathbf{y}_i^T\left(\mathbf{I} + N_i \mathrm{E}\left[\mathbf{V}^T\mathbf{W}\mathbf{V}\right]\right)\mathbf{y}_i + \mathrm{const} \tag{33}$$

Therefore $q^*(\mathbf{Y})$ is a product of Gaussian distributions.

$$q^*(\mathbf{Y}) = \prod_{i=1}^{M} \mathcal{N}\left(\mathbf{y}_i | \overline{\mathbf{y}}_i, \mathbf{L}_{\mathbf{y}_i}^{-1}\right) \tag{34}$$

$$\mathbf{L}_{\mathbf{y}_i} = \mathbf{I} + N_i \mathrm{E}\left[\mathbf{V}^T \mathbf{W} \mathbf{V}\right] \tag{35}$$

$$\overline{\mathbf{y}}_i = \mathbf{L}_{\mathbf{y}_i}^{-1} \mathrm{E}\left[\mathbf{V}^T \mathbf{W}\left(\mathbf{F}_i - N_i \mu\right)\right] \tag{36}$$

$$= \mathbf{L}_{\mathbf{y}_i}^{-1}\left(\mathrm{E}\left[\mathbf{V}\right]^T \mathrm{E}\left[\mathbf{W}\right] \mathbf{F}_i - N_i \mathrm{E}\left[\mathbf{V}^T \mathbf{W} \mu\right]\right) \tag{37}$$

The optimum for $q^*(\tilde{\mathbf{v}}'_r)$:

$$\ln q^*(\tilde{\mathbf{v}}'_r) = \mathrm{E}_{\mathbf{Y},\mathbf{W},\alpha,\tilde{\mathbf{v}}'_{s\neq r}}\left[\ln P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha)\right] + \text{const} \tag{38}$$

$$= \mathrm{E}_{\mathbf{Y},\mathbf{W},\tilde{\mathbf{v}}'_{s\neq r}}\left[\ln P(\mathbf{\Phi}|\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W})\right] + \mathrm{E}_{\alpha,\mathbf{v}'_{s\neq r}}\left[\ln P(\mathbf{V}|\alpha)\right] + \mathrm{E}_{\mu_{s\neq r}}\left[\ln P(\mu)\right] + \text{const} \tag{39}$$

$$= -\frac{1}{2}\sum_{i=1}^{M} \mathrm{tr}\left(\mathrm{E}\left[\mathbf{W}\right]\left(-2\mathbf{F}_i \mathrm{E}\left[\tilde{\mathbf{y}}_i\right]^T \mathrm{E}_{\tilde{\mathbf{v}}'_{s\neq r}}\left[\tilde{\mathbf{V}}\right]^T + N_i \mathrm{E}_{\tilde{\mathbf{v}}'_{s\neq r}}\left[\tilde{\mathbf{V}} \mathrm{E}\left[\tilde{\mathbf{y}}_i \tilde{\mathbf{y}}_i^T\right] \tilde{\mathbf{V}}^T\right]\right)\right)$$
$$- \frac{1}{2}\sum_{q=1}^{n_y} \mathrm{E}\left[\alpha_q\right] \mathrm{E}_{\mathbf{v}'_{s\neq r}}\left[\mathbf{v}_q^T \mathbf{v}_q\right] - \frac{1}{2}\beta_r \left(\mu_r - \mu_{0_r}\right)^2 + \text{const} \tag{40}$$

$$= -\frac{1}{2}\mathrm{tr}\left(\mathrm{E}\left[\mathbf{W}\right]\left(-2\mathbf{C}\mathrm{E}_{\tilde{\mathbf{v}}'_{s\neq r}}\left[\tilde{\mathbf{V}}\right]^T + \mathrm{E}_{\tilde{\mathbf{v}}'_{s\neq r}}\left[\tilde{\mathbf{V}} \mathbf{R}_{\tilde{\mathbf{y}}} \tilde{\mathbf{V}}^T\right]\right)\right)$$
$$- \frac{1}{2}\sum_{q=1}^{n_y} \mathrm{E}\left[\alpha_q\right] \mathrm{E}_{\mathbf{v}'_{s\neq r}}\left[\mathbf{v}_q^T \mathbf{v}_q\right] - \frac{1}{2}\beta_r \left(\mu_r - \mu_{0_r}\right)^2 + \text{const} \tag{41}$$

$$= -\frac{1}{2}\mathrm{tr}\left(-2\mathrm{E}_{\tilde{\mathbf{v}}'_{s\neq r}}\left[\tilde{\mathbf{V}}\right]^T \mathrm{E}\left[\mathbf{W}\right] \mathbf{C} + \mathrm{E}_{\tilde{\mathbf{v}}'_{s\neq r}}\left[\tilde{\mathbf{V}}^T \mathrm{E}\left[\mathbf{W}\right] \tilde{\mathbf{V}}\right] \mathbf{R}_{\tilde{\mathbf{y}}}\right)$$
$$- \frac{1}{2}\mathbf{v}_r'^T \mathrm{diag}\left(\mathrm{E}\left[\alpha\right]\right) \mathbf{v}'_r - \frac{1}{2}\beta_r \left(\mu_r - \mu_{0_r}\right)^2 + \text{const} \tag{42}$$

$$= -\frac{1}{2}\mathrm{tr}\left(-2\sum_{s=1}^{d} \tilde{\mathbf{v}}'_r \overline{w}_{rs} \mathbf{C}_s + 2\sum_{s\neq r} \tilde{\mathbf{v}}'_r \overline{w}_{rs} \mathrm{E}\left[\tilde{\mathbf{v}}'_s\right]^T \mathbf{R}_{\tilde{\mathbf{y}}} + \tilde{\mathbf{v}}'_r \overline{w}_{rr} \tilde{\mathbf{v}}_r'^T \mathbf{R}_{\tilde{\mathbf{y}}}\right)$$
$$- \frac{1}{2}\mathbf{v}_r'^T \mathrm{diag}\left(\mathrm{E}\left[\alpha\right]\right) \mathbf{v}'_r - \frac{1}{2}\beta_r \left(\mu_r - \mu_{0_r}\right)^2 + \text{const} \tag{43}$$

$$= -\frac{1}{2}\mathrm{tr}\left(-2\tilde{\mathbf{v}}'_r\left(\overline{w}_{rr}\mathbf{C}_r + \sum_{s\neq r} \overline{w}_{rs}\left(\mathbf{C}_s - \mathrm{E}\left[\tilde{\mathbf{v}}'_s\right]^T \mathbf{R}_{\tilde{\mathbf{y}}}\right)\right) + \tilde{\mathbf{v}}'_r \tilde{\mathbf{v}}_r'^T \overline{w}_{rr} \mathbf{R}_{\tilde{\mathbf{y}}}\right)$$
$$- \frac{1}{2}\mathbf{v}_r'^T \mathrm{diag}\left(\mathrm{E}\left[\alpha\right]\right) \mathbf{v}'_r - \frac{1}{2}\beta_r \left(\mu_r - \mu_{0_r}\right)^2 + \text{const} \tag{44}$$

$$= -\frac{1}{2}\mathrm{tr}\left(-2\tilde{\mathbf{v}}'_r\left(\overline{w}_{rr}\mathbf{C}_r + \sum_{s\neq r} \overline{w}_{rs}\left(\mathbf{C}_s - \mathrm{E}\left[\tilde{\mathbf{v}}'_s\right]^T \mathbf{R}_{\tilde{\mathbf{y}}}\right)\right) + \tilde{\mathbf{v}}'_r \tilde{\mathbf{v}}_r'^T \overline{w}_{rr} \mathbf{R}_{\tilde{\mathbf{y}}}\right)$$
$$- \frac{1}{2}\tilde{\mathbf{v}}_r'^T \mathrm{diag}\left(\overline{\tilde{\alpha}}_r\right) \tilde{\mathbf{v}}'_r + \beta_r \mu_r \mu_{0_r} + \text{const} \tag{45}$$

$$= -\frac{1}{2}\mathrm{tr}\left(-2\tilde{\mathbf{v}}'_r\left(\overline{w}_{rr}\mathbf{C}_r + \sum_{s\neq r} \overline{w}_{rs}\left(\mathbf{C}_s - \mathrm{E}\left[\tilde{\mathbf{v}}'_s\right]^T \mathbf{R}_{\tilde{\mathbf{y}}}\right) + \beta_r \tilde{\mu}_{0_r}^T\right)\right.$$
$$\left. + \tilde{\mathbf{v}}'_r \tilde{\mathbf{v}}_r'^T \left(\mathrm{diag}\left(\overline{\tilde{\alpha}}_r\right) + \overline{w}_{rr} \mathbf{R}_{\tilde{\mathbf{y}}}\right)\right) \tag{46}$$

where $\overline{w}_{rs}$ is the element r, s of $\mathrm{E}[\mathbf{W}]$,

$$\mathbf{C} = \sum_{i=1}^{M} \mathbf{F}_i \mathrm{E}[\tilde{\mathbf{y}}_i]^T \tag{47}$$

$$\mathbf{R}_{\tilde{\mathbf{y}}} = \sum_{i=1}^{M} N_i \mathrm{E}\left[\tilde{\mathbf{y}}_i \tilde{\mathbf{y}}_i^T\right] \tag{48}$$

$$\mathrm{E}\left[\tilde{\mathbf{y}}_i \tilde{\mathbf{y}}_i^T\right] = \begin{bmatrix} \mathrm{E}\left[\mathbf{y}_i \mathbf{y}_i^T\right] & \mathrm{E}[\mathbf{y}_i] \\ \mathrm{E}[\mathbf{y}_i]^T & 1 \end{bmatrix} \tag{49}$$

$$\overline{\tilde{\alpha}}_r = \begin{bmatrix} \mathrm{E}[\alpha] \\ \beta_r \end{bmatrix} \qquad \tilde{\mu}_{0_r} = \begin{bmatrix} \mathbf{0}_{n_y \times 1} \\ \mu_{0_r} \end{bmatrix} \tag{50}$$

and $\mathbf{C}_r$ is the r^{th} row of $\mathbf{C}$.

Then $q^*(\tilde{\mathbf{v}}'_r)$ is a Gaussian distribution:

$$q^*(\tilde{\mathbf{v}}'_r) = \mathcal{N}\left(\tilde{\mathbf{v}}'_r | \overline{\tilde{\mathbf{v}}}'_r, \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r}\right) \tag{51}$$

$$\mathbf{L}_{\tilde{\mathbf{V}}_r} = \mathrm{diag}\left(\overline{\tilde{\alpha}}_r\right) + \overline{w}_{rr} \mathbf{R}_{\tilde{\mathbf{y}}} \tag{52}$$

$$\overline{\tilde{\mathbf{v}}}'_r = \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r} \left(\overline{w}_{rr} \mathbf{C}_r^T + \sum_{s \neq r} \overline{w}_{rs} \left(\mathbf{C}_s^T - \mathbf{R}_{\tilde{\mathbf{y}}} \overline{\tilde{\mathbf{v}}}'_s \right) + \beta_r \tilde{\mu}_{0_r} \right) \tag{53}$$

The optimum for $q^*(\alpha)$:

$$\ln q^*(\alpha) = \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha)\right] + \text{const} \tag{54}$$

$$= \mathrm{E}_{\mathbf{V}}\left[\ln P(\mathbf{V}|\alpha)\right] + \ln P(\alpha | a_\alpha, b_\alpha) + \text{const} \tag{55}$$

$$= \sum_{q=1}^{n_y} \frac{d}{2} \ln \alpha_q - \frac{1}{2} \alpha_q \mathrm{E}\left[\mathbf{v}_q^T \mathbf{v}_q\right] + (a_\alpha - 1) \ln \alpha_q - b_\alpha \alpha_q + \text{const} \tag{56}$$

$$= \sum_{q=1}^{n_y} \left(\frac{d}{2} + a_\alpha - 1 \right) \ln \alpha_q - \alpha_q \left(b_\alpha + \frac{1}{2} \mathrm{E}\left[\mathbf{v}_q^T \mathbf{v}_q\right] \right) + \text{const} \tag{57}$$

$$\tag{58}$$

Then $q^*(\alpha)$ is a product of Gammas:

$$q^*(\alpha) = \prod_{q=1}^{n_y} \mathcal{G}\left(\alpha_q | a'_\alpha, b'_{\alpha_q}\right) \tag{59}$$

$$a'_\alpha = a_\alpha + \frac{d}{2} \tag{60}$$

$$b'_{\alpha_q} = b_\alpha + \frac{1}{2} \mathrm{E}\left[\mathbf{v}_q^T \mathbf{v}_q\right] \tag{61}$$

The optimum for $q^*(\mathbf{W})$ in the non-informative case:

$$\ln q^*(\mathbf{W}) = \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\alpha}\left[\ln P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha)\right] + \text{const} \tag{62}$$

$$= \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V}}\left[\ln P(\mathbf{\Phi}|\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W})\right] + \ln P(\mathbf{W}) + \text{const} \tag{63}$$

$$= \frac{N}{2} \ln |\mathbf{W}| - \frac{d+1}{2} \ln |\mathbf{W}| - \frac{1}{2} \mathrm{tr}(\mathbf{W}\mathbf{K}) + \text{const} \tag{64}$$

where

$$\mathbf{K} = \sum_{i=1}^{M} \mathrm{E}\left[\mathbf{S}_i - \mathbf{F}_i \tilde{\mathbf{y}}_i^T \tilde{\mathbf{V}}^T - \tilde{\mathbf{V}} \tilde{\mathbf{y}}_i \mathbf{F}_i^T + N_i \tilde{\mathbf{V}} \tilde{\mathbf{y}}_i \tilde{\mathbf{y}}_i^T \tilde{\mathbf{V}}^T\right] \tag{65}$$

$$= \mathbf{S} - \mathbf{C}\mathrm{E}\left[\tilde{\mathbf{V}}\right]^T - \mathrm{E}\left[\tilde{\mathbf{V}}\right] \mathbf{C}^T + \mathrm{E}_{\tilde{\mathbf{V}}}\left[\tilde{\mathbf{V}} \mathbf{R}_{\tilde{\mathbf{y}}} \tilde{\mathbf{V}}^T\right] \tag{66}$$

Then $q^*(\mathbf{W})$ is Wishart distributed:

$$P(\mathbf{W}) = \mathcal{W}(\mathbf{W}|\mathbf{\Psi}, \nu) \quad \text{if } \nu > d \tag{67}$$
$$\mathbf{\Psi}^{-1} = \mathbf{K} \tag{68}$$
$$\nu = N \tag{69}$$

The optimum for $q^*(\mathbf{W})$ in the informative case:

$$\ln q^*(\mathbf{W}) = \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\alpha}[\ln P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha)] + \text{const} \tag{70}$$
$$= \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V}}[\ln P(\mathbf{\Phi}|\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W})] + \ln P(\mathbf{W}) + \text{const} \tag{71}$$
$$= \frac{N}{2}\ln|\mathbf{W}| + \frac{\nu_0 - d - 1}{2}\ln|\mathbf{W}| - \frac{1}{2}\mathrm{tr}\left(\mathbf{W}\left(\mathbf{\Psi}_0^{-1} + \mathbf{K}\right)\right) + \text{const} \tag{72}$$

Then $q^*(\mathbf{W})$ is Wishart distributed:

$$P(\mathbf{W}) = \mathcal{W}(\mathbf{W}|\mathbf{\Psi}, \nu) \tag{73}$$
$$\mathbf{\Psi}^{-1} = \mathbf{\Psi}_0^{-1} + \mathbf{K} \tag{74}$$
$$\nu = \nu_0 + N \tag{75}$$

Finally, we evaluate the expectations:

$$\mathrm{E}\left[\alpha_q\right] = \frac{a'_\alpha}{b'_{\alpha_q}} \tag{76}$$

$$\overline{\tilde{\mathbf{V}}} = \mathrm{E}\left[\tilde{\mathbf{V}}\right] = \begin{bmatrix} \overline{\tilde{\mathbf{v}}}_1'^T \\ \overline{\tilde{\mathbf{v}}}_2'^T \\ \vdots \\ \overline{\tilde{\mathbf{v}}}_d'^T \end{bmatrix} \tag{77}$$

$$\overline{\mathbf{W}} = \mathrm{E}\left[\mathbf{W}\right] = \nu\mathbf{\Psi} \tag{78}$$

$$\mathrm{E}\left[\mathbf{v}_q^T\mathbf{v}_q\right] = \sum_{r=1}^{d}\mathrm{E}\left[\mathbf{v}_{rq}'^T\mathbf{v}'_{rq}\right] \tag{79}$$

$$= \sum_{r=1}^{d}\mathbf{L}^{-1}_{\tilde{\mathbf{V}}_{rqq}} + \overline{\mathbf{v}}'^2_{rq} \tag{80}$$

$$\mathrm{E}\left[\mathbf{V}^T\mathbf{W}\mathbf{V}\right] = \mathrm{E}\left[\mathbf{V}'\overline{\mathbf{W}}\mathbf{V}'^T\right] \tag{81}$$

$$= \sum_{r=1}^{d}\sum_{s=1}^{d}\overline{w}_{rs}\mathrm{E}\left[\mathbf{v}'_r\mathbf{v}_s'^T\right] \tag{82}$$

$$= \sum_{r=1}^{d}\overline{w}_{rr}\mathbf{\Sigma}_{\mathbf{V}_r} + \sum_{r=1}^{d}\sum_{s=1}^{d}\overline{w}_{rs}\overline{\mathbf{v}}'_r\overline{\mathbf{v}}_s'^T \tag{83}$$

$$= \sum_{r=1}^{d}\overline{w}_{rr}\mathbf{\Sigma}_{\mathbf{V}_r} + \overline{\mathbf{V}}^T\overline{\mathbf{W}\mathbf{V}} \tag{84}$$

$$\mathrm{E}\left[\mathbf{V}^T\mathbf{W}\mu\right] = \sum_{r=1}^{d}\overline{w}_{rr}\mathbf{\Sigma}_{\mathbf{V}\mu_r} + \sum_{r=1}^{d}\sum_{s=1}^{d}\overline{w}_{rs}\overline{\mathbf{v}}'_r\overline{\mu}_s \tag{85}$$

$$= \sum_{r=1}^{d}\overline{w}_{rr}\mathbf{\Sigma}_{\mathbf{V}\mu_r} + \overline{\mathbf{V}}^T\overline{\mathbf{W}}\overline{\mu} \tag{86}$$

$$\mathrm{E}\left[\tilde{\mathbf{V}}\mathbf{R}_{\tilde{\mathbf{y}}}\tilde{\mathbf{V}}^T\right] = \sum_{r=1}^{n_y}\sum_{s=1}^{n_y}r_{\tilde{\mathbf{y}}_{rs}}\mathrm{E}\left[\tilde{\mathbf{v}}_r\tilde{\mathbf{v}}_s^T\right] \tag{87}$$

$$= \sum_{r=1}^{n_y}\sum_{s=1}^{n_y}r_{\tilde{\mathbf{y}}_{rs}}\left[\mathrm{E}\left[\tilde{v}_{r_i}\tilde{v}_{s_j}\right]\right]_{d\times d} \tag{88}$$

$$= \sum_{r=1}^{n_y}\sum_{s=1}^{n_y}r_{\tilde{\mathbf{y}}_{rs}}\left[\mathrm{E}\left[\tilde{v}'_{i_r}\tilde{v}'_{j_s}\right]\right]_{d\times d} \tag{89}$$

$$= \sum_{r=1}^{n_y}\sum_{s=1}^{n_y}r_{\tilde{\mathbf{y}}_{rs}}\left[\mathrm{E}\left[\tilde{v}'_{i_r}\right]\mathrm{E}\left[\tilde{v}'_{j_s}\right]\right]_{d\times d} + \sum_{r=1}^{n_y}\sum_{s=1}^{n_y}r_{\tilde{\mathbf{y}}_{rs}}\mathrm{diag}\left(\left[\sigma_{\tilde{\mathbf{V}}_{i_{rs}}}\right]_d\right) \tag{90}$$

$$= \sum_{r=1}^{n_y}\sum_{s=1}^{n_y}r_{\tilde{\mathbf{y}}_{rs}}\left[\mathrm{E}\left[\tilde{v}_{r_i}\right]\mathrm{E}\left[\tilde{v}_{s_j}\right]\right]_{d\times d} + \mathrm{diag}\left(\rho\right) \tag{91}$$

$$= \sum_{r=1}^{n_y}\sum_{s=1}^{n_y}r_{\tilde{\mathbf{y}}_{rs}}\mathrm{E}\left[\tilde{\mathbf{v}}_r\right]\mathrm{E}\left[\tilde{\mathbf{v}}_s\right]^T + \mathrm{diag}\left(\rho\right) \tag{92}$$

$$= \overline{\tilde{\mathbf{V}}}\mathbf{R}_{\tilde{\mathbf{y}}}\overline{\tilde{\mathbf{V}}}^T + \mathrm{diag}\left(\rho\right) \tag{93}$$

where $\tilde{v}_{r_i}$ is the i^{th} element of $\tilde{\mathbf{v}}_r$, $\tilde{v}'_{i_r}$ is the r^{th} element of $\tilde{\mathbf{v}}'_i$,

$$\boldsymbol{\Sigma}_{\tilde{\mathbf{V}}_r} = \begin{bmatrix} \boldsymbol{\Sigma}_{\mathbf{V}_r} & \boldsymbol{\Sigma}_{\mathbf{V}\mu_r} \\ \boldsymbol{\Sigma}^T_{\mathbf{V}\mu_r} & \boldsymbol{\Sigma}_{\mu_r} \end{bmatrix} = \left[\sigma_{\tilde{\mathbf{V}}_{r_{ij}}}\right]_{n_y \times n_y} = \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r} \tag{94}$$

$$\rho = \begin{bmatrix} \rho_1 & \rho_2 & \dots & \rho_d \end{bmatrix}^T \tag{95}$$

$$\rho_i = \sum_{r=1}^{n_y} \sum_{s=1}^{n_y} \left(\mathbf{R}_{\tilde{\mathbf{y}}} \circ \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_i}\right)_{rs} \tag{96}$$

and $\circ$ is the Hadamard product.

5.2.1 Distributions with deterministic annealing

If we use annealing, for a parameter κ, we have:

$$q^*(\mathbf{Y}) = \prod_{i=1}^{M} \mathcal{N}\left(\mathbf{y}_i | \overline{\mathbf{y}}_i, 1/\kappa\, \mathbf{L}^{-1}_{\mathbf{y}_i}\right) \tag{97}$$

$$q^*(\tilde{\mathbf{v}}'_r) = \mathcal{N}\left(\tilde{\mathbf{v}}'_r | \overline{\tilde{\mathbf{v}}}'_r, 1/\kappa\, \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r}\right) \tag{98}$$

$$q^*(\mathbf{W}) = \mathcal{W}(\mathbf{W} | 1/\kappa\, \boldsymbol{\Psi}, \kappa(\nu - d - 1) + d + 1) \quad \text{if } \kappa(\nu - d - 1) + 1 > 0 \tag{99}$$

$$q^*(\alpha) = \prod_{q=1}^{n_y} \mathcal{G}\left(\alpha_q | a'_\alpha, b'_{\alpha_q}\right) \tag{100}$$

$$a'_\alpha = \kappa\left(a_\alpha + \frac{d}{2} - 1\right) + 1 \tag{101}$$

$$b'_{\alpha_q} = \kappa\left(b_\alpha + \frac{1}{2}\mathrm{E}\left[\mathbf{v}_q^T \mathbf{v}_q\right]\right)\ . \tag{102}$$

5.3 Variational lower bound

The lower bound is given by

$$\begin{aligned} \mathcal{L} =& \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P(\boldsymbol{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W})\right] + \mathrm{E}_{\mathbf{Y}}\left[\ln P(\mathbf{Y})\right] + \mathrm{E}_{\mathbf{V},\alpha}\left[\ln P(\mathbf{V}|\alpha)\right] \\ &+ \mathrm{E}_{\alpha}\left[\ln P(\alpha)\right] + \mathrm{E}_{\mu}\left[\ln P(\mu)\right] + \mathrm{E}_{\mathbf{W}}\left[\ln P(\mathbf{W})\right] \\ &- \mathrm{E}_{\mathbf{Y}}\left[\ln q(\mathbf{Y})\right] - \mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln q\left(\tilde{\mathbf{V}}\right)\right] - \mathrm{E}_{\alpha}\left[\ln q(\alpha)\right] - \mathrm{E}_{\mathbf{W}}\left[\ln q(\mathbf{W})\right] \end{aligned} \tag{103}$$

The term $\mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P(\boldsymbol{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W})\right]$:

$$\begin{aligned} \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P(\boldsymbol{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W})\right] =& \frac{N}{2}\mathrm{E}\left[\ln|\mathbf{W}|\right] - \frac{Nd}{2}\ln(2\pi) \\ &- \frac{1}{2}\mathrm{tr}\left(\overline{\mathbf{W}}\left(\mathbf{S} - 2\mathbf{C}\overline{\tilde{\mathbf{V}}}^T + \mathrm{E}\left[\tilde{\mathbf{V}}\mathbf{R}_{\tilde{\mathbf{y}}}\tilde{\mathbf{V}}^T\right]\right)\right) \end{aligned} \tag{104}$$

$$\begin{aligned} =& \frac{N}{2}\ln\overline{\mathbf{W}} - \frac{Nd}{2}\ln(2\pi) - \frac{1}{2}\mathrm{tr}\left(\overline{\mathbf{W}}\mathbf{S}\right) \\ &- \frac{1}{2}\mathrm{tr}\left(-2\overline{\tilde{\mathbf{V}}}^T\overline{\mathbf{W}}\mathbf{C} + \mathrm{E}\left[\tilde{\mathbf{V}}^T\mathbf{W}\tilde{\mathbf{V}}\right]\mathbf{R}_{\tilde{\mathbf{y}}}\right) \end{aligned} \tag{105}$$

where

$$\ln\overline{\mathbf{W}} = \mathrm{E}\left[\ln|\mathbf{W}|\right] \tag{106}$$

$$= \sum_{i=1}^{d}\psi\left(\frac{\nu + 1 - i}{2}\right) + d\ln 2 + \ln|\boldsymbol{\Psi}| \tag{107}$$

and ψ is the digamma function.

The term $\mathrm{E}_{\mathbf{Y}}\left[\ln P\left(\mathbf{Y}\right)\right]$:

$$\mathrm{E}_{\mathbf{Y}}\left[\ln P\left(\mathbf{Y}\right)\right] = -\frac{Mn_y}{2}\ln(2\pi) - \frac{1}{2}\mathrm{tr}\left(\sum_{i=1}^{M}\mathrm{E}\left[\mathbf{y}_i\mathbf{y}_i^T\right]\right) \tag{108}$$

$$= -\frac{Mn_y}{2}\ln(2\pi) - \frac{1}{2}\mathrm{tr}\left(\mathrm{P}\right) \tag{109}$$

where

$$\mathrm{P} = \sum_{i=1}^{M}\mathrm{E}\left[\mathbf{y}_i\mathbf{y}_i^T\right] \tag{110}$$

The term $\mathrm{E}_{\mathbf{V},\alpha}\left[\ln P\left(\mathbf{V}|\alpha\right)\right]$:

$$\mathrm{E}_{\mathbf{V},\alpha}\left[\ln P\left(\mathbf{V}|\alpha\right)\right] = -\frac{n_y d}{2}\ln(2\pi) + \frac{d}{2}\sum_{q=1}^{n_y}\mathrm{E}\left[\ln\alpha_q\right] - \frac{1}{2}\sum_{q=1}^{n_y}\mathrm{E}\left[\alpha_q\right]\mathrm{E}\left[\mathbf{v}_q^T\mathbf{v}_q\right] \tag{111}$$

where

$$\mathrm{E}\left[\ln\alpha_q\right] = \psi(a'_\alpha) - \ln b'_{\alpha_q}\ . \tag{112}$$

The term $\mathrm{E}_{\alpha}\left[\ln P\left(\alpha\right)\right]$:

$$\mathrm{E}_{\alpha}\left[\ln P\left(\alpha\right)\right] = n_y\left(a_\alpha\ln b_\alpha - \ln\Gamma\left(a_\alpha\right)\right) + \sum_{q=1}^{n_y}(a_\alpha - 1)\mathrm{E}\left[\ln\alpha_q\right] - b_\alpha\mathrm{E}\left[\alpha_q\right] \tag{113}$$

$$= n_y\left(a_\alpha\ln b_\alpha - \ln\Gamma\left(a_\alpha\right)\right) + (a_\alpha - 1)\sum_{q=1}^{n_y}\mathrm{E}\left[\ln\alpha_q\right] - b_\alpha\sum_{q=1}^{n_y}\mathrm{E}\left[\alpha_q\right] \tag{114}$$

The term $\mathrm{E}_{\mu}\left[\ln P\left(\mu\right)\right]$:

$$\mathrm{E}_{\mu}\left[\ln P\left(\mu\right)\right] = -\frac{d}{2}\ln(2\pi) + \frac{1}{2}\sum_{r=1}^{d}\ln\beta_r - \frac{1}{2}\sum_{r=1}^{d}\beta_r\left(\mathrm{E}\left[\mu_r^2\right] - 2\mu_{0_r}\mathrm{E}\left[\mu_r\right] + \mu_{0_r}^2\right) \tag{115}$$

$$= -\frac{d}{2}\ln(2\pi) + \frac{1}{2}\sum_{r=1}^{d}\ln\beta_r - \frac{1}{2}\sum_{r=1}^{d}\beta_r\left(\mathbf{\Sigma}_{\mu_r} + \mathrm{E}\left[\mu_r\right]^2 - 2\mu_{0_r}\mathrm{E}\left[\mu_r\right] + \mu_{0_r}^2\right) \tag{116}$$

The term $\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right]$ for the non-informative case:

$$\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right] = -\frac{d+1}{2}\ln\overline{\mathbf{W}}\ . \tag{117}$$

The term $\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right]$ for the informative case:

$$\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right] = \ln B\left(\mathbf{\Psi}_0,\nu_0\right) + \frac{\nu_0 - d - 1}{2}\ln\overline{\mathbf{W}} - \frac{\nu}{2}\mathrm{tr}\left(\mathbf{\Psi}_0^{-1}\mathbf{\Psi}\right) \tag{118}$$

The term $\mathrm{E}_{\mathbf{Y}}\left[\ln q\left(\mathbf{Y}\right)\right]$:

$$\mathrm{E}_{\mathbf{Y}}\left[\ln q\left(\mathbf{Y}\right)\right] = -\frac{Mn_y}{2}\ln(2\pi) + \frac{1}{2}\sum_{i=1}^{M}\ln\left|\mathbf{L}_{\mathbf{y}_i}\right| - \frac{1}{2}\mathrm{tr}\left(\mathbf{L}_{\mathbf{y}_i}\mathrm{E}\left[\left(\mathbf{y}_i-\overline{\mathbf{y}}_i\right)\left(\mathbf{y}_i-\overline{\mathbf{y}}_i\right)^T\right]\right) \tag{119}$$

$$= -\frac{Mn_y}{2}\ln(2\pi) + \frac{1}{2}\sum_{i=1}^{M}\ln\left|\mathbf{L}_{\mathbf{y}_i}\right| - \frac{1}{2}\sum_{i=1}^{M}\mathrm{tr}\left(\mathbf{L}_{\mathbf{y}_i}\left(\mathrm{E}\left[\mathbf{y}_i\mathbf{y}_i^T\right] - \overline{\mathbf{y}}_i\mathrm{E}\left[\mathbf{y}_i\right]^T - \mathrm{E}\left[\mathbf{y}_i\right]\overline{\mathbf{y}}_i^T + \overline{\mathbf{y}}_i\overline{\mathbf{y}}_i^T\right)\right) \tag{120}$$

$$= -\frac{Mn_y}{2}\ln(2\pi) + \frac{1}{2}\sum_{i=1}^{M}\ln\left|\mathbf{L}_{\mathbf{y}_i}\right| - \frac{1}{2}\sum_{i=1}^{M}\mathrm{tr}\left(\mathbf{I}\right) \tag{121}$$

$$= -\frac{Mn_y}{2}(\ln(2\pi)+1) + \frac{1}{2}\sum_{i=1}^{M}\ln\left|\mathbf{L}_{\mathbf{y}_i}\right| \tag{122}$$

The term $\mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln q\left(\tilde{\mathbf{V}}\right)\right]$:

$$\mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln q\left(\tilde{\mathbf{V}}\right)\right] = -\frac{d(n_y+1)}{2}\left(\ln(2\pi)+1\right) + \frac{1}{2}\sum_{r=1}^{d}\ln\left|\mathbf{L}_{\tilde{\mathbf{V}}_r}\right| \tag{123}$$

The term $\mathrm{E}_{\alpha}\left[\ln q\left(\alpha\right)\right]$:

$$\mathrm{E}_{\alpha}\left[\ln q\left(\alpha\right)\right] = -\sum_{q=1}^{n_y}\mathrm{H}\left[q\left(\alpha_q\right)\right] \tag{124}$$

$$= \sum_{q=1}^{n_y}(a'_\alpha - 1)\psi(a'_\alpha) + \ln b'_{\alpha_q} - a'_\alpha - \ln\Gamma\left(a'_\alpha\right) \tag{125}$$

$$= n_y\left((a'_\alpha - 1)\psi(a'_\alpha) - a'_\alpha - \ln\Gamma\left(a'_\alpha\right)\right) + \sum_{q=1}^{n_y}\ln b'_{\alpha_q} \tag{126}$$

The term $\mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right]$:

$$\mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right] = -\mathrm{H}\left[q\left(\mathbf{W}\right)\right] \tag{127}$$

$$= \ln B\left(\mathbf{\Psi},\nu\right) + \frac{\nu-d-1}{2}\ln\overline{\mathbf{W}} - \frac{\nu d}{2} \tag{128}$$

where

$$B(\mathbf{A},N) = \frac{1}{2^{Nd/2}Z_{Nd}}\left|\mathbf{A}\right|^{-N/2} \tag{129}$$

$$Z_{Nd} = \pi^{d(d-1)/4}\prod_{i=1}^{d}\Gamma\left((N+1-i)/2\right) \tag{130}$$

5.4 Hyperparameter optimization

We can set the Hyperparameters $(\mu_0, \beta, a_\alpha, b_\alpha)$ manually or estimate them from the development data maximizing the lower bound.

we derive for a_α

$$\frac{\partial\mathcal{L}}{\partial a_\alpha} = n_y\left(\ln b_\alpha - \psi(a_\alpha)\right) + \sum_{q=1}^{n_y}\mathrm{E}\left[\ln\alpha_q\right] = 0 \quad \Longrightarrow \tag{131}$$

$$\psi(a_\alpha) = \ln b_\alpha + \frac{1}{n_y}\sum_{q=1}^{n_y}\mathrm{E}\left[\ln\alpha_q\right] \tag{132}$$

We derive for b_α:

$$\frac{\partial \mathcal{L}}{\partial b_\alpha} = \frac{n_y a_\alpha}{b_\alpha} - \sum_{q=1}^{n_y} \mathrm{E}\left[\alpha_q\right] = \mathbf{0} \quad \Longrightarrow \tag{133}$$

$$b_\alpha = \left(\frac{1}{n_y a_\alpha} \sum_{q=1}^{n_y} \mathrm{E}\left[\alpha_q\right] \right)^{-1} \tag{134}$$

We solve these equations with the procedure described in [5]. We write

$$\psi(a) = \ln b + c \tag{135}$$

$$b = \frac{a}{d} \tag{136}$$

where

$$c = \frac{1}{n_y} \sum_{q=1}^{n_y} \mathrm{E}\left[\ln \alpha_q\right] \tag{137}$$

$$d = \frac{1}{n_y} \sum_{q=1}^{n_y} \mathrm{E}\left[\alpha_q\right] \tag{138}$$

Then

$$f(a) = \psi(a) - \ln a + \ln d - c = 0 \tag{139}$$

We can solve for a using Newton-Rhaphson iterations:

$$a_{new} = a - \frac{f(a)}{f'(a)} = \tag{140}$$

$$= a \left(1 - \frac{\psi(a) - \ln a + \ln d - c}{a\psi'(a) - 1} \right) \tag{141}$$

This algorithm does not assure that a remains positive. We can put a minimum value for a. Alternatively we can solve the equation for $\tilde{a}$ such as $a = exp(\tilde{a})$.

$$\tilde{a}_{new} = \tilde{a} - \frac{f(\tilde{a})}{f'(\tilde{a})} = \tag{142}$$

$$= \tilde{a} - \frac{\psi(a) - \ln a + \ln d - c}{\psi'(a)a - 1} \tag{143}$$

Taking exponential in both sides:

$$a_{new} = a \exp\left(- \frac{\psi(a) - \ln a + \ln d - c}{\psi'(a)a - 1} \right) \tag{144}$$

We derive for μ_0:

$$\frac{\partial \mathcal{L}}{\partial \mu_0} = \mathbf{0} \quad \Longrightarrow \tag{145}$$

$$\mu_0 = \mathrm{E}\left[\mu\right] \tag{146}$$

We derive for β:

$$\frac{\partial \mathcal{L}}{\partial \beta} = \mathbf{0} \quad \Longrightarrow \tag{147}$$

$$\beta_r^{-1} = \mathbf{\Sigma}_{\mu_r} + \mathrm{E}\left[\mu_r\right]^2 - 2\mu_{0_r}\mathrm{E}\left[\mu_r\right] + \mu_{0_r}^2 \tag{148}$$

If we take an isotropic prior for μ:

$$\beta^{-1} = \frac{1}{d} \sum_{r=1}^{d} \mathbf{\Sigma}_{\mu_r} + \mathrm{E}\left[\mu_r\right]^2 - 2\mu_{0_r}\mathrm{E}\left[\mu_r\right] + \mu_{0_r}^2 \tag{149}$$

5.5 Minimum divergence

We assume a more general prior for the hidden variables:

$$P(\mathbf{y}) = \mathcal{N}\left(\mathbf{y}|\mu_{\mathbf{y}}, \mathbf{\Lambda}_{\mathbf{y}}^{-1}\right) \tag{150}$$

To minimize the divergence we maximize the part of $\mathcal{L}$ that depends on $\mu_{\mathbf{y}}$:

$$\mathcal{L}(\mu_{\mathbf{y}}, \mathbf{\Lambda}_{\mathbf{y}}) = \sum_{i=1}^{M} \mathrm{E}_{\mathbf{Y}}\left[\ln \mathcal{N}\left(\mathbf{y}|\mu_{\mathbf{y}}, \mathbf{\Lambda}_{\mathbf{y}}^{-1}\right)\right] \tag{151}$$

The, we get

$$\mu_{\mathbf{y}} = \frac{1}{M}\sum_{i=1}^{M} \mathrm{E}_{\mathbf{Y}}\left[\mathbf{y}_i\right] \tag{152}$$

$$\mathbf{\Sigma}_{\mathbf{y}} = \mathbf{\Lambda}_{\mathbf{y}}^{-1} = \frac{1}{M}\sum_{i=1}^{M} \mathrm{E}_{\mathbf{Y}}\left[\left(\mathbf{y}_i - \mu_{\mathbf{y}}\right)\left(\mathbf{y}_i - \mu_{\mathbf{y}}\right)^T\right] \tag{153}$$

$$= \frac{1}{M}\sum_{i=1}^{M} \mathrm{E}_{\mathbf{Y}}\left[\mathbf{y}_i\mathbf{y}_i^T\right] - \mu_{\mathbf{y}}\mu_{\mathbf{y}}^T \tag{154}$$

We have a transform $\mathbf{y} = \phi(\mathbf{y}')$ such as $\mathbf{y}'$ has a standard prior:

$$\mathbf{y} = \mu_{\mathbf{y}} + (\mathbf{\Sigma}_{\mathbf{y}}^{1/2})^T\mathbf{y}' \tag{155}$$

we also can write that as

$$\tilde{\mathbf{y}} = \mathbf{J}\tilde{\mathbf{y}}' \tag{156}$$

where

$$\mathbf{J} = \begin{bmatrix} (\mathbf{\Sigma}_{\mathbf{y}}^{1/2})^T & \mu_{\mathbf{y}} \\ \mathbf{0}^T & 1 \end{bmatrix} \tag{157}$$

Now, we get $q(\tilde{\mathbf{v}}'_r)$ such us if we apply the transform $\mathbf{y}' = \phi^{-1}(\mathbf{y})$, the term $\mathrm{E}\left[\ln P(\mathbf{X}|\mathbf{Y}, \mathbf{W})\right]$ of $\mathcal{L}$ remains constant:

$$\overline{\tilde{\mathbf{v}}}'_r \leftarrow \mathbf{J}^T\overline{\tilde{\mathbf{v}}}'_r \tag{158}$$

$$\mathbf{L}_{\tilde{\mathbf{V}}_r}^{-1} \leftarrow \mathbf{J}^T\mathbf{L}_{\tilde{\mathbf{V}}_r}^{-1}\mathbf{J} \tag{159}$$

$$\mathbf{L}_{\tilde{\mathbf{V}}_r} \leftarrow \mathbf{G}^T\mathbf{L}_{\tilde{\mathbf{V}}_r}\mathbf{G} \tag{160}$$

where

$$\mathbf{G} = \left(\mathbf{J}^T\right)^{-1} \tag{161}$$

$$= \begin{bmatrix} (\mathbf{\Sigma}_{\mathbf{y}}^{1/2})^{-1} & \mathbf{0} \\ -\mu_{\mathbf{y}}^T(\mathbf{\Sigma}_{\mathbf{y}}^{1/2})^{-1} & 1 \end{bmatrix} \tag{162}$$

6 Variational inference with Gaussian-Gamma priors for V, Gaussian for μ and Gamma for W

6.1 Model priors

In section 5, we saw that if we use a full covariance $\mathbf{W}$ we had a full covariance posterior for $\tilde{\mathbf{V}}$. Then, to get a tractable solution, we forced independence between the the rows of $\tilde{\mathbf{V}}$ when choosing the variational partition function.

In this section, we are going to assume that we have applied a rotation to the data such as we can consider that $\mathbf{W}$ is going to remain diagonal during the VB iteration.

Then we are going to place a broad Gamma prior over each element of the diagonal of $\mathbf{W}$:

$$P(\mathbf{W}) = \prod_{r=1}^{d} \mathcal{G}(w_{rr}|a_w, b_w) \tag{163}$$

We also consider the case of an isotropic $\mathbf{W}$ ($\mathbf{W} = w\mathbf{I}$). Then the prior is

$$P(w) = \mathcal{G}(w|a_w, b_w) \tag{164}$$

6.2 Variational distributions

We write the joint distribution of the latent variables:

$$\begin{aligned} P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha|\mu_0, \beta, a_\alpha, b_\alpha, a_w, b_w) =& P(\mathbf{\Phi}|\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}) P(\mathbf{Y}) P(\mathbf{V}|\alpha) \\ & P(\alpha|a, b) P(\mu|\mu_0, \beta) P(\mathbf{W}|a_w, b_w) \end{aligned} \tag{165}$$

Following, the conditioning on $\mu_0, \beta, a_\alpha, b_\alpha, a_w, b_w$ will be dropped for convenience.

Now, we consider the partition of the posterior:

$$P(\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha|\mathbf{\Phi}) \approx q(\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha) = q(\mathbf{Y}) q\left(\tilde{\mathbf{V}}\right) q(\mathbf{W}) q(\alpha) \tag{166}$$

The optimum for $q^*(\mathbf{Y})$ and $q^*(\alpha)$ are the same as in section 5.2.

The optimum for $q^*\left(\tilde{\mathbf{V}}\right)$:

$$\ln q^*\left(\tilde{\mathbf{V}}\right) = \mathrm{E}_{\mathbf{Y},\mathbf{W},\alpha}\left[\ln P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha)\right] + \text{const} \tag{167}$$

$$= \mathrm{E}_{\mathbf{Y},\mathbf{W}}\left[\ln P(\mathbf{\Phi}|\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W})\right] + \mathrm{E}_\alpha\left[\ln P(\mathbf{V}|\alpha)\right] + \ln P(\mu) + \text{const} \tag{168}$$

$$\begin{aligned} =& -\frac{1}{2}\mathrm{tr}\left(-2\tilde{\mathbf{V}}^T \mathrm{E}[\mathbf{W}] \mathbf{C} + \tilde{\mathbf{V}}^T \mathrm{E}[\mathbf{W}] \tilde{\mathbf{V}} \mathbf{R}_{\tilde{\mathbf{y}}}\right) \\ & - \frac{1}{2}\sum_{r=1}^{d} \mathbf{v}_r'^T \mathrm{diag}(\mathrm{E}[\alpha]) \mathbf{v}_r' - \frac{1}{2}\sum_{r=1}^{d} \beta_r (\mu_r - \mu_{0_r})^2 + \text{const} \end{aligned} \tag{169}$$

$$= -\frac{1}{2}\sum_{r=1}^{d} \mathrm{tr}\left(-2\tilde{\mathbf{v}}_r'\left(\overline{w}_{rr}\mathbf{C}_r + \beta_r \tilde{\mu}_{0_r}^T\right) + \tilde{\mathbf{v}}_r'\tilde{\mathbf{v}}_r'^T\left(\mathrm{diag}\left(\overline{\widetilde{\alpha}}_r\right) + \overline{w}_{rr}\mathbf{R}_{\tilde{\mathbf{y}}}\right)\right) \tag{170}$$

Then $q^*\left(\tilde{\mathbf{V}}\right)$ is a product of Gaussian distributions:

$$q^*\left(\tilde{\mathbf{V}}\right) = \prod_{r=1}^{d} \mathcal{N}\left(\tilde{\mathbf{v}}_r'|\overline{\widetilde{\mathbf{w}}}_r', \mathbf{L}_{\tilde{\mathbf{V}}_r}^{-1}\right) \tag{171}$$

$$\mathbf{L}_{\tilde{\mathbf{V}}_r} = \mathrm{diag}\left(\overline{\widetilde{\alpha}}_r\right) + \overline{w}_{rr}\mathbf{R}_{\tilde{\mathbf{y}}} \tag{172}$$

$$\overline{\widetilde{\mathbf{v}}}_r' = \mathbf{L}_{\tilde{\mathbf{V}}_r}^{-1}\left(\overline{w}_{rr}\mathbf{C}_r^T + \beta_r \tilde{\mu}_{0_r}\right) \tag{173}$$

The optimum for $q^*(\mathbf{W})$:

$$\ln q^*(\mathbf{W}) = \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\alpha}\left[\ln P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}, \alpha)\right] + \text{const} \tag{174}$$

$$= \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V}}\left[\ln P(\mathbf{\Phi}|\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W})\right] + \ln P(\mathbf{W}) + \text{const} \tag{175}$$

$$= \sum_{r=1}^{d} \frac{N}{2} \ln w_{rr} - \frac{1}{2} w_{rr} k_{rr} + (a_w - 1) \ln w_{rr} - b_w w_{rr} + \text{const} \tag{176}$$

$$= \sum_{r=1}^{d} \left(a_w + \frac{N}{2} - 1\right) \ln w_{rr} - \left(b_w + \frac{1}{2} k_{rr}\right) w_{rr} + \text{const} \tag{177}$$

where

$$\mathbf{K} = \text{diag}\left(\mathbf{S} - \mathbf{C}\text{E}\left[\tilde{\mathbf{V}}\right]^T - \text{E}\left[\tilde{\mathbf{V}}\right]\mathbf{C}^T + \text{E}_{\tilde{\mathbf{V}}}\left[\tilde{\mathbf{V}}\mathbf{R}_{\tilde{\mathbf{y}}}\tilde{\mathbf{V}}^T\right]\right) \tag{178}$$

Then $q^*(\mathbf{W})$ is a product of Gammas:

$$q^*(\mathbf{W}) = \prod_{r=1}^{d} \mathcal{G}\left(w_{rr}|a'_w, b'_{w_r}\right) \tag{179}$$

$$a'_w = a_w + \frac{N}{2} \tag{180}$$

$$b'_{w_r} = b_w + \frac{1}{2}k_{rr} \tag{181}$$

If we force an isotropic $\mathbf{W}$, the optimum $q^*(\mathbf{W})$ is

$$\ln q^*(\mathbf{W}) = \frac{Nd}{2}\ln w - \frac{1}{2}wk + (a_w - 1)\ln w - b_w w + \text{const} \tag{182}$$

$$= \left(a_w + \frac{Nd}{2} - 1\right)\ln w - \left(b_w + \frac{1}{2}k\right)w + \text{const} \tag{183}$$

where

$$k = \text{tr}\left(\mathbf{S} - \mathbf{C}\text{E}\left[\tilde{\mathbf{V}}\right]^T - \text{E}\left[\tilde{\mathbf{V}}\right]\mathbf{C}^T + \text{E}_{\tilde{\mathbf{V}}}\left[\tilde{\mathbf{V}}\mathbf{R}_{\tilde{\mathbf{y}}}\tilde{\mathbf{V}}^T\right]\right) \tag{184}$$

$$= \text{tr}\left(\mathbf{S} - 2\mathbf{C}\text{E}\left[\tilde{\mathbf{V}}\right]^T\right) + \text{tr}\left(\text{E}\left[\tilde{\mathbf{V}}^T\tilde{\mathbf{V}}\right]\mathbf{R}_{\tilde{\mathbf{y}}}\right) \tag{185}$$

Then $q^*(\mathbf{W})$ is a Gamma distribution:

$$q^*(\mathbf{W}) = \mathcal{G}\left(w|a'_w, b'_w\right) \tag{186}$$

$$a'_w = a_w + \frac{Nd}{2} \tag{187}$$

$$b'_w = b_w + \frac{1}{2}k \tag{188}$$

Finally, we evaluate the expectations:

$$\text{E}\left[\tilde{\mathbf{V}}^T\tilde{\mathbf{V}}\right] = \text{E}\left[\tilde{\mathbf{V}}'\tilde{\mathbf{V}}'^T\right] \tag{189}$$

$$= \sum_{r=1}^{d}\mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r} + \text{E}\left[\tilde{\mathbf{V}}'\right]\text{E}\left[\tilde{\mathbf{V}}'\right]^T \tag{190}$$

$$= \sum_{r=1}^{d}\mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r} + \text{E}\left[\tilde{\mathbf{V}}\right]^T\text{E}\left[\tilde{\mathbf{V}}\right] \tag{191}$$

6.3 Variational lower bound

The lower bound is given by

$$\begin{aligned}\mathcal{L} =& \text{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P\left(\mathbf{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W}\right)\right] + \text{E}_{\mathbf{Y}}\left[\ln P\left(\mathbf{Y}\right)\right] + \text{E}_{\mathbf{V},\alpha}\left[\ln P\left(\mathbf{V}|\alpha\right)\right] \\ &+ \text{E}_{\alpha}\left[\ln P\left(\alpha\right)\right] + \text{E}_{\mu}\left[\ln P\left(\mu\right)\right] + \text{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right] \\ &- \text{E}_{\mathbf{Y}}\left[\ln q\left(\mathbf{Y}\right)\right] - \text{E}_{\tilde{\mathbf{V}}}\left[\ln q\left(\tilde{\mathbf{V}}\right)\right] - \text{E}_{\alpha}\left[\ln q\left(\alpha\right)\right] - \text{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right]\end{aligned} \tag{192}$$

The term $\text{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P\left(\mathbf{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W}\right)\right]$:

$$\begin{aligned}\text{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P\left(\mathbf{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W}\right)\right] =& \frac{N}{2}\sum_{r=1}^{d}\ln\overline{w}_{rr} - \frac{Nd}{2}\ln(2\pi) - \frac{1}{2}\text{tr}\left(\overline{\mathbf{W}}\mathbf{S}\right) \\ &- \frac{1}{2}\text{tr}\left(-2\overline{\tilde{\mathbf{V}}}^T\overline{\mathbf{W}}\mathbf{C} + \text{E}\left[\tilde{\mathbf{V}}^T\mathbf{W}\tilde{\mathbf{V}}\right]\mathbf{R}_{\tilde{\mathbf{y}}}\right)\end{aligned} \tag{193}$$

where

$$\ln \overline{w}_{rr} = \mathrm{E}\left[\ln |w_{rr}|\right] \tag{194}$$
$$= \psi(a'_w) - \ln b'_{w_r} \tag{195}$$

and ψ is the digamma function.

The term $\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right]$ with non-isotropic $\mathbf{W}$:

$$\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right] = \sum_{r=1}^{d} a_w \ln b_w + \left(a_w - 1\right) \mathrm{E}\left[\ln w_{rr}\right] - b_w \mathrm{E}\left[w_{rr}\right] - \ln \Gamma\left(a_w\right) \tag{196}$$
$$= d\left(a_w \ln b_w - \ln \Gamma\left(a_w\right)\right) + \left(a_w - 1\right) \sum_{r=1}^{d} \mathrm{E}\left[\ln w_{rr}\right] - b_w \sum_{r=1}^{d} \mathrm{E}\left[w_{rr}\right] \tag{197}$$

The term $\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right]$ with isotropic $\mathbf{W}$:

$$\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right] = a_w \ln b_w - \ln \Gamma\left(a_w\right) + \left(a_w - 1\right) \mathrm{E}\left[\ln w\right] - b_w \mathrm{E}\left[w\right] \tag{198}$$

The term $\mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right]$ with non-isotropic $\mathbf{W}$:

$$\mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right] = -\sum_{r=1}^{d} \mathrm{H}\left[q\left(w_{rr}\right)\right] \tag{199}$$
$$= d\left(\left(a'_w - 1\right)\psi(a'_w) - a'_w - \ln \Gamma\left(a'_w\right)\right) + \sum_{r=1}^{d} \ln b'_{w_r} \tag{200}$$

The term $\mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right]$ with isotropic $\mathbf{W}$:

$$\mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right] = -\mathrm{H}\left[q\left(w\right)\right] \tag{201}$$
$$= \left(a'_w - 1\right)\psi(a'_w) - a'_w - \ln \Gamma\left(a'_w\right) + \ln b'_w \tag{202}$$

The rest of terms are the same as in section 6.3.

6.4 Hyperparameter optimization

We can estimate the parameters (a_w, b_w) from the development data maximizing the lower bound.

For non-isotropic $\mathbf{W}$:

we derive for a_w

$$\frac{\partial \mathcal{L}}{\partial a_w} = d\left(\ln b_w - \psi(a_w)\right) + \sum_{r=1}^{d} \mathrm{E}\left[\ln w_{rr}\right] = 0 \quad \Longrightarrow \tag{203}$$

$$\psi(a_w) = \ln b_w + \frac{1}{d} \sum_{r=1}^{d} \mathrm{E}\left[\ln w_{rr}\right] \tag{204}$$

We derive for b_w:

$$\frac{\partial \mathcal{L}}{\partial b_w} = \frac{d a_w}{b} - \sum_{r=1}^{d} \mathrm{E}\left[w_{rr}\right] = \mathbf{0} \quad \Longrightarrow \tag{205}$$

$$b_w = \left(\frac{1}{d a_w} \sum_{r=1}^{d} \mathrm{E}\left[w_{rr}\right]\right)^{-1} \tag{206}$$

For isotropic $\mathbf{W}$:

we derive for a_w

$$\frac{\partial \mathcal{L}}{\partial a_w} = \ln b_w - \psi(a_w) + \mathrm{E}\left[\ln w\right] = 0 \quad \Longrightarrow \tag{207}$$

$$\psi(a_w) = \ln b_w + \mathrm{E}\left[\ln w\right] \tag{208}$$

We derive for b_w:

$$\frac{\partial \mathcal{L}}{\partial b_w} = \frac{a_w}{b} - \mathrm{E}\left[w\right] = \mathbf{0} \quad \Longrightarrow \tag{209}$$

$$b_w = \left(\frac{1}{a_w}\mathrm{E}\left[w\right]\right)^{-1} \tag{210}$$

We can solve these equations by Newton-Rhapson iterations as described in section 5.4.

7 Variational inference with full covariance Gaussian prior for V and μ and Wishart for W

7.1 Model priors

Lets assume that we compute the posterior of model parameters given a development database with a large amount of data. If we want to compute the model posterior for a small database we could use the posterior given the large database as prior.

Thus, we take a prior distribution for $\tilde{\mathbf{V}}$

$$P\left(\tilde{\mathbf{V}}\right) = \prod_{r=1}^{d} \mathcal{N}\left(\mathbf{v}'_r | \overline{\mathbf{v}}'_{0r}, \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_{0r}}\right) \tag{211}$$

The prior for $\mathbf{W}$ is

$$P\left(\mathbf{W}\right) = \mathcal{W}\left(\mathbf{W} | \mathbf{\Psi}_0, \nu_0\right) \tag{212}$$

The parameters $\overline{\mathbf{v}}'_{0r}$, $\mathbf{L}^{-1}_{\tilde{\mathbf{V}}_{0r}}$, Ψ_0 and ν_0 are computed with the large dataset.

7.2 Variational distributions

The joint distribution of the latent variables:

$$P\left(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}\right) = P\left(\mathbf{\Phi} | \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}\right) P\left(\mathbf{Y}\right) P\left(\tilde{\mathbf{V}}\right) P\left(\mathbf{W}\right) \tag{213}$$

Now, we consider the partition of the posterior:

$$P\left(\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W} | \mathbf{\Phi}\right) \approx q\left(\mathbf{Y}, \tilde{\mathbf{V}}, \mathbf{W}\right) = q\left(\mathbf{Y}\right) \prod_{r=1}^{d} q\left(\tilde{\mathbf{v}}'_r\right) q\left(\mathbf{W}\right) \tag{214}$$

The optimum for $q^*\left(\mathbf{Y}\right)$ is the same as in section 5.2.
The optimum for $q^*\left(\tilde{\mathbf{v}}'_r\right)$:

$$\ln q^*\left(\tilde{\mathbf{v}}'_r\right) = \mathrm{E}_{\mathbf{Y},\mathbf{W},\tilde{\mathbf{v}}'_{s\neq r}}\left[\ln P\left(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}\right)\right] + \mathrm{const} \tag{215}$$

$$= \mathrm{E}_{\mathbf{Y},\mathbf{W},\tilde{\mathbf{v}}'_{s\neq r}}\left[\ln P\left(\mathbf{\Phi} | \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}\right)\right] + \ln P\left(\tilde{\mathbf{V}}\right) + \mathrm{const} \tag{216}$$

$$\begin{aligned} &= -\frac{1}{2}\mathrm{tr}\left(-2\tilde{\mathbf{v}}'_r\left(\overline{w}_{rr}\mathbf{C}_r + \sum_{s\neq r}\overline{w}_{rs}\left(\mathbf{C}_s - \mathrm{E}\left[\tilde{\mathbf{v}}'_s\right]^T \mathbf{R}_{\tilde{\mathbf{y}}}\right)\right) + \tilde{\mathbf{v}}'_r\tilde{\mathbf{v}}'^T_r\overline{w}_{rr}\mathbf{R}_{\tilde{\mathbf{y}}}\right) \\ &\quad - \frac{1}{2}(\mathbf{v}'_r - \overline{\mathbf{v}}'_{0r})^T\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}(\mathbf{v}'_r - \overline{\mathbf{v}}'_{0r}) + \mathrm{const} \end{aligned} \tag{217}$$

$$\begin{aligned} &= -\frac{1}{2}\mathrm{tr}\left(-2\tilde{\mathbf{v}}'_r\left(\overline{\mathbf{v}}'^T_{0r}\mathbf{L}_{\tilde{\mathbf{V}}_{0r}} + \overline{w}_{rr}\mathbf{C}_r + \sum_{s\neq r}\overline{w}_{rs}\left(\mathbf{C}_s - \mathrm{E}\left[\tilde{\mathbf{v}}'_s\right]^T \mathbf{R}_{\tilde{\mathbf{y}}}\right)\right)\right. \\ &\quad \left. + \tilde{\mathbf{v}}'_r\tilde{\mathbf{v}}'^T_r\left(\mathbf{L}_{\tilde{\mathbf{V}}_{0r}} + \overline{w}_{rr}\mathbf{R}_{\tilde{\mathbf{y}}}\right)\right) \end{aligned} \tag{218}$$

Then $q^*\left(\tilde{\mathbf{v}}'_r\right)$ is a Gaussian distribution:

$$q^*\left(\tilde{\mathbf{v}}'_r\right) = \mathcal{N}\left(\tilde{\mathbf{v}}'_r | \overline{\tilde{\mathbf{v}}}'_r, \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r}\right) \tag{219}$$

$$\mathbf{L}_{\tilde{\mathbf{V}}_r} = \mathbf{L}_{\tilde{\mathbf{V}}_{0r}} + \overline{w}_{rr}\mathbf{R}_{\tilde{\mathbf{y}}} \tag{220}$$

$$\overline{\tilde{\mathbf{v}}}'_r = \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r}\left(\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\overline{\mathbf{v}}'_{0r} + \overline{w}_{rr}\mathbf{C}_r^T + \sum_{s\neq r}\overline{w}_{rs}\left(\mathbf{C}_s^T - \mathbf{R}_{\tilde{\mathbf{y}}}\overline{\tilde{\mathbf{v}}}'_s\right)\right) \tag{221}$$

The optimum for $q^*\left(\mathbf{W}\right)$:

$$\ln q^*\left(\mathbf{W}\right) = \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\alpha}\left[\ln P\left(\mathbf{\Phi},\mathbf{Y},\mu,\mathbf{V},\mathbf{W},\alpha\right)\right] + \text{const} \tag{222}$$

$$= \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V}}\left[\ln P\left(\mathbf{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W}\right)\right] + \ln P\left(\mathbf{W}\right) + \text{const} \tag{223}$$

$$= \frac{N}{2}\ln|\mathbf{W}| + \frac{\nu_0 - d - 1}{2}\ln|\mathbf{W}| - \frac{1}{2}\mathrm{tr}\left(\mathbf{W}\left(\mathbf{\Psi}_0^{-1} + \mathbf{K}\right)\right) + \text{const} \tag{224}$$

where

$$\mathbf{K} = \mathbf{S} - \mathbf{C}\mathrm{E}\left[\tilde{\mathbf{V}}\right]^T - \mathrm{E}\left[\tilde{\mathbf{V}}\right]\mathbf{C}^T + \mathrm{E}_{\tilde{\mathbf{V}}}\left[\tilde{\mathbf{V}}\mathbf{R}_{\tilde{\mathbf{y}}}\tilde{\mathbf{V}}^T\right] \tag{225}$$

Then $q^*\left(\mathbf{W}\right)$ is Wishart distributed:

$$P\left(\mathbf{W}\right) = \mathcal{W}\left(\mathbf{W}|\mathbf{\Psi},\nu\right) \tag{226}$$

$$\mathbf{\Psi}^{-1} = \mathbf{\Psi}_0^{-1} + \mathbf{K} \tag{227}$$

$$\nu = \nu_0 + N \tag{228}$$

7.2.1 Distributions with deterministic annealing

If we use annealing, for a parameter κ, we have:

$$q^*\left(\mathbf{W}\right) = \mathcal{W}\left(\mathbf{W}|1/\kappa\ \mathbf{\Psi}, \kappa(\nu_0 + N - d - 1) + d + 1\right) \quad \text{if } \kappa(\nu_0 + N - d - 1) + 1 > 0 \tag{229}$$

7.3 Variational lower bound

The lower bound is given by

$$\begin{aligned}\mathcal{L} =& \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P\left(\mathbf{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W}\right)\right] + \mathrm{E}_{\mathbf{Y}}\left[\ln P\left(\mathbf{Y}\right)\right] + \mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln P\left(\tilde{\mathbf{V}}\right)\right] + \mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right] \\ & - \mathrm{E}_{\mathbf{Y}}\left[\ln q\left(\mathbf{Y}\right)\right] - \mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln q\left(\tilde{\mathbf{V}}\right)\right] - \mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right]\end{aligned} \tag{230}$$

The term $\mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln P\left(\tilde{\mathbf{V}}\right)\right]$:

$$\begin{aligned}
\mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln P\left(\tilde{\mathbf{V}}\right)\right] =& -\frac{n_y d}{2}\ln(2\pi) + \frac{1}{2}\sum_{r=1}^{d}\ln\left|\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\right| \\
& -\frac{1}{2}\sum_{r=1}^{d}\mathrm{tr}\left(\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\mathrm{E}\left[\left(\mathbf{v}'_r-\overline{\mathbf{v}}'_{0r}\right)\left(\mathbf{v}'_r-\overline{\mathbf{v}}'_{0r}\right)^T\right]\right) && (231)\\
=& -\frac{n_y d}{2}\ln(2\pi) + \frac{1}{2}\sum_{r=1}^{d}\ln\left|\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\right| \\
& -\frac{1}{2}\sum_{r=1}^{d}\mathrm{tr}\left(\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\left(\mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r}+\overline{\mathbf{v}}'_r\overline{\mathbf{v}}'^T_r-\overline{\mathbf{v}}'_{0r}\overline{\mathbf{v}}'^T_r-\overline{\mathbf{v}}'_r\overline{\mathbf{v}}'^T_{0r}+\overline{\mathbf{v}}'_{0r}\overline{\mathbf{v}}'^T_{0r}\right)\right) && (232)\\
=& -\frac{n_y d}{2}\ln(2\pi) + \frac{1}{2}\sum_{r=1}^{d}\ln\left|\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\right| \\
& -\frac{1}{2}\sum_{r=1}^{d}\mathrm{tr}\left(\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r}\right)-\frac{1}{2}\sum_{r=1}^{d}(\overline{\mathbf{v}}'_r-\overline{\mathbf{v}}'_{0r})^T\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}(\overline{\mathbf{v}}'_r-\overline{\mathbf{v}}'_{0r}) && (233)\\
=& -\frac{n_y d}{2}\ln(2\pi) + \frac{1}{2}\sum_{r=1}^{d}\ln\left|\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\right| \\
& -\frac{1}{2}\sum_{r=1}^{d}\mathrm{tr}\left(\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r}\right)-\frac{1}{2}\sum_{r=1}^{d}\mathrm{tr}\left(\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\left(\overline{\mathbf{v}}'_r-\overline{\mathbf{v}}'_{0r}\right)\left(\overline{\mathbf{v}}'_r-\overline{\mathbf{v}}'_{0r}\right)^T\right) && (234)
\end{aligned}$$

The term $\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right]$:

$$\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right] = \ln B\left(\mathbf{\Psi}_0,\nu_0\right)+\frac{\nu_0-d-1}{2}\ln\overline{\mathbf{W}}-\frac{\nu}{2}\mathrm{tr}\left(\mathbf{\Psi}_0^{-1}\mathbf{\Psi}\right) \tag{235}$$

where

$$\begin{aligned}
\ln\overline{\mathbf{W}} =& \mathrm{E}\left[\ln|\mathbf{W}|\right] && (236)\\
=& \sum_{i=1}^{d}\psi\left(\frac{\nu+1-i}{2}\right)+d\ln 2+\ln|\mathbf{\Psi}| && (237)
\end{aligned}$$

and ψ is the digamma function.

The term $\mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right]$

$$\begin{aligned}
\mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right] =& -\mathrm{H}\left[q\left(\mathbf{W}\right)\right] && (238)\\
=& \ln B\left(\mathbf{\Psi},\nu\right)+\frac{\nu-d-1}{2}\ln\overline{\mathbf{W}}-\frac{\nu d}{2} && (239)
\end{aligned}$$

The rest of terms are the same as the ones in section 5.3.

8 Variational inference with full covariance Gaussian prior for V and μ and Gamma for W

8.1 Model priors

Thus, we take a prior distribution for $\tilde{\mathbf{V}}$

$$P\left(\tilde{\mathbf{V}}\right)=\prod_{r=1}^{d}\mathcal{N}\left(\mathbf{v}'_r|\overline{\mathbf{v}}'_{0r},\mathbf{L}^{-1}_{\tilde{\mathbf{V}}_{0r}}\right) \tag{240}$$

The prior for non-isotropic $\mathbf{W}$ is

$$P(\mathbf{W}) = \prod_{r=1}^{d} \mathcal{G}(w_{rr}|a_w, b_{w_r}) \tag{241}$$

The prior for isotropic $\mathbf{W}$ is

$$P(\mathbf{W}) = \mathcal{G}(w|a_w, b_w) \tag{242}$$

The parameters $\overline{\mathbf{v}}'_{0r}$, $\mathbf{L}^{-1}_{\tilde{\mathbf{V}}_{0r}}$, a_w and b_w are computed with the large dataset.

8.2 Variational distributions

The joint distribution of the latent variables:

$$P(\mathbf{\Phi}, \mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}) = P(\mathbf{\Phi}|\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}) P(\mathbf{Y}) P\left(\tilde{\mathbf{V}}\right) P(\mathbf{W}) \tag{243}$$

Now, we consider the partition of the posterior:

$$P(\mathbf{Y}, \mu, \mathbf{V}, \mathbf{W}|\mathbf{\Phi}) \approx q\left(\mathbf{Y}, \tilde{\mathbf{V}}, \mathbf{W}\right) = q(\mathbf{Y}) q\left(\tilde{\mathbf{V}}\right) q(\mathbf{W}) \tag{244}$$

The optimum for $q^*(\mathbf{Y})$ is the same as in section 5.2.
Then optimum for $q^*\left(\tilde{\mathbf{V}}\right)$ is:

$$q^*\left(\tilde{\mathbf{V}}\right) = \prod_{r=1}^{d} \mathcal{N}\left(\tilde{\mathbf{v}}'_r|\overline{\tilde{\mathbf{v}}}'_r, \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r}\right) \tag{245}$$

$$\mathbf{L}_{\tilde{\mathbf{V}}_r} = \mathbf{L}_{\tilde{\mathbf{V}}_{0r}} + \overline{w}_{rr}\mathbf{R}_{\tilde{\mathbf{y}}} \tag{246}$$

$$\overline{\tilde{\mathbf{v}}}'_r = \mathbf{L}^{-1}_{\tilde{\mathbf{V}}_r}\left(\mathbf{L}_{\tilde{\mathbf{V}}_{0r}}\overline{\mathbf{v}}'_{0r} + \overline{w}_{rr}\mathbf{C}_r^T\right) \tag{247}$$

The optimum for $q^*(\mathbf{W})$ for non-isotropic $\mathbf{W}$:

$$q^*(\mathbf{W}) = \prod_{r=1}^{d} \mathcal{G}\left(w_{rr}|a'_w, b'_{w_r}\right) \tag{248}$$

$$a'_w = a_w + \frac{N}{2} \tag{249}$$

$$b'_{w_r} = b_{w_r} + \frac{1}{2}k_{rr} \tag{250}$$

where

$$\mathbf{K} = \text{diag}\left(\mathbf{S} - \mathbf{C}\text{E}\left[\tilde{\mathbf{V}}\right]^T - \text{E}\left[\tilde{\mathbf{V}}\right]\mathbf{C}^T + \text{E}_{\tilde{\mathbf{V}}}\left[\tilde{\mathbf{V}}\mathbf{R}_{\tilde{\mathbf{y}}}\tilde{\mathbf{V}}^T\right]\right) \tag{251}$$

The optimum for $q^*(\mathbf{W})$ for isotropic $\mathbf{W}$:

$$q^*(\mathbf{W}) = \mathcal{G}\left(w|a'_w, b'_w\right) \tag{252}$$

$$a'_w = a_w + \frac{Nd}{2} \tag{253}$$

$$b'_w = b_w + \frac{1}{2}k \tag{254}$$

where

$$k = \text{tr}\left(\mathbf{S} - 2\mathbf{C}\text{E}\left[\tilde{\mathbf{V}}\right]^T\right) + \text{tr}\left(\text{E}\left[\tilde{\mathbf{V}}^T\tilde{\mathbf{V}}\right]\mathbf{R}_{\tilde{\mathbf{y}}}\right) \tag{255}$$

8.3 Variational lower bound

The lower bound is given by

$$\begin{aligned}\mathcal{L} =& \mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P\left(\mathbf{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W}\right)\right] + \mathrm{E}_{\mathbf{Y}}\left[\ln P\left(\mathbf{Y}\right)\right] + \mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln P\left(\tilde{\mathbf{V}}\right)\right] + \mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right] \\ & - \mathrm{E}_{\mathbf{Y}}\left[\ln q\left(\mathbf{Y}\right)\right] - \mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln q\left(\tilde{\mathbf{V}}\right)\right] - \mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right] \end{aligned} \tag{256}$$

The term $\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right]$ with non-isotropic $\mathbf{W}$:

$$\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right] = \sum_{r=1}^{d} a_w \ln b_{w_r} + \left(a_w - 1\right)\mathrm{E}\left[\ln w_{rr}\right] - b_{w_r}\mathrm{E}\left[w_{rr}\right] - \ln\Gamma\left(a_w\right) \tag{257}$$

$$= -\, d\ln\Gamma\left(a_w\right) + a_w\sum_{r=1}^{d}\ln b_{w_r} + \left(a_w - 1\right)\sum_{r=1}^{d}\mathrm{E}\left[\ln w_{rr}\right] - \sum_{r=1}^{d} b_{w_r}\mathrm{E}\left[w_{rr}\right] \tag{258}$$

The terms $\mathrm{E}_{\mathbf{Y}}\left[\ln P\left(\mathbf{Y}\right)\right]$, $\mathrm{E}_{\mathbf{Y}}\left[\ln q\left(\mathbf{Y}\right)\right]$ and $\mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln q\left(\tilde{\mathbf{V}}\right)\right]$ are the same as in section 5.3.

The terms $\mathrm{E}_{\mathbf{Y},\mu,\mathbf{V},\mathbf{W}}\left[\ln P\left(\mathbf{\Phi}|\mathbf{Y},\mu,\mathbf{V},\mathbf{W}\right)\right]$, $\mathrm{E}_{\mathbf{W}}\left[\ln P\left(\mathbf{W}\right)\right]$ with isotropic $\mathbf{W}$ and $\mathrm{E}_{\mathbf{W}}\left[\ln q\left(\mathbf{W}\right)\right]$ are the same as is in section 6.3.

The term $\mathrm{E}_{\tilde{\mathbf{V}}}\left[\ln P\left(\tilde{\mathbf{V}}\right)\right]$ is the same as in section 7.3.

References

[1] Jesús Villalba, "SPLDA," Tech. Rep., University of Zaragoza, Zaragoza, 2011.

[2] Jesús Villalba and Niko Brummer, "Towards Fully Bayesian Speaker Recognition: Integrating Out the Between-Speaker Covariance," in *Interspeech 2011*, Florence, 2011, pp. 28–31.

[3] C M Bishop, "Variational principal components," *9th International Conference on Artificial Neural Networks ICANN 99*, vol. 1, no. 470, pp. 509–514, 1999.

[4] Jesús Villalba, "Fully Bayesian Two-Covariance Model," Tech. Rep., University of Zaragoza, Zaragoza (Spain), 2010.

[5] Matthew J Beal, "Variational algorithms for approximate Bayesian inference," *Philosophy*, vol. 38, no. May, pp. 1–281, 2003.